

GEOLOGICAL SURVEY OF ALABAMA

Berry H. (Nick) Tew, Jr.
State Geologist

ECOSYSTEM INVESTIGATIONS PROGRAM

Patrick E. O'Neil
Director

**RESULTS OF A SURVEY OF THE MUSSEL FAUNA AT
SELECTED STATIONS IN THE BLACK WARRIOR RIVER
SYSTEM, ALABAMA, 2009-10**

OPEN-FILE REPORT 1006

By
Stuart W. McGregor¹, E. Anne Wynn¹ and Jeffrey T. Garner²

¹Geological Survey of Alabama

²Alabama Department of Conservation and Natural Resources
Division of Wildlife and Freshwater Fisheries

In cooperation with the Alabama Department of Conservation and Natural Resources
Division of Wildlife and Freshwater Fisheries
Project 127

Tuscaloosa, Alabama
2010

CONTENTS

	Page
Abstract.....	1
Introduction.....	1
Acknowledgments.....	3
Study area.....	3
Methods.....	3
Species accounts	3
Results and discussion	5
Recommendations.....	16
References cited	17

FIGURES

Figure 1. Map of the study area showing mussel sampling locations.....	4
--	---

TABLES

Table 1. Summary information for stations sampled in the Black Warrior River system, Alabama, 2009-10.....	6
Table 2. Overview of freshwater mussels collected in the Black Warrior River, Alabama, 2009	8
Table 3. Overview of freshwater mussels collected in tributaries of the Black Warrior River, Alabama, 2009-10	9
Table 4. Comparison of freshwater mussel species in Locust Fork and tributaries reported by Hartfield (1990) and the current study	12

RESULTS OF A SURVEY OF THE MUSSEL FAUNA AT SELECTED STATIONS IN THE BLACK WARRIOR RIVER SYSTEM, ALABAMA, 2009-10

By

Stuart W. McGregor, E. Anne Wynn, and Jeffrey T. Garner

ABSTRACT

During the summers of 2009 and 2010, sampling for mussels in the Black Warrior River system yielded 21 species from 16 main channel and 43 tributary stations. A cumulative total of about 8.0 hours bottom time was spent sampling main channel stations (average 0.5 hour per station) and about 58 hours in tributaries (average 0.7 hour per station). Main channel collections yielded 675 live or fresh dead individuals. The most numerically dominant and frequently encountered species were *Plectomerus dombeyanus*, the Bankclimber (9 stations, 28.4 collected per hour), *Quadrula apiculata*, the Southern Mapleleaf (11 stations, 23.8 collected per hour), and *Obliquaria reflexa*, the Threehorn Wartyback (11 stations, 22.0 collected per hour). The federally listed threatened species *Potamilus inflatus*, Inflated Heelsplitter, was collected live at two main channel stations. Most tributary stations yielded no evidence of mussels or only weathered dead shells. Some notable exceptions were a few Locust Fork and Blackburn Fork stations, where seven species were found live. No federally listed or State of Alabama Conservation Priority species were found live or fresh dead in tributaries, but weathered dead valves of *Elliptio arctata*, Delicate Spike, a Highest Conservation Priority species, were found in Locust Fork, and a badly eroded valve of what may be *Pleurobema rubellum*, the Warrior Pigtoe, was collected from Davis Creek.

INTRODUCTION

The mussel and fish faunas of the Mobile River Basin are noteworthy for their high degrees of endemism and diversity. Those phenomena can be attributed to the large size of the basin, numerous habitat types available due to the diverse physiography of the basin, geographic barriers such as the Fall Line, and the proximity of the basin to adjacent drainages with diverse faunas (Williams, 1982). Hinkley (1906) reported 40 mussel species from the Tombigbee River system alone, while Williams and others (1992) reported that 50 species were known to have occurred in the upper Tombigbee River system (upstream of the confluence of the Tombigbee and Black Warrior Rivers) and 48 in the Black Warrior system, adjusted to taxonomic revisions in recent decades. Williams and others (2008), in a comprehensive review of the mussels of

Alabama, tallied 51 species known from the Black Warrior system. However, effects of anthropogenic factors such as impoundment, eutrophication, sedimentation, pollution, and channel modifications and resultant population fragmentation of mussels and their obligate fish hosts, have suppressed gene flow and caused a steep and possibly fatal decline in the fauna (Hartfield, 1994; Mott and Hartfield, 1994). While the mussel population in Alabama remains among the most diverse on Earth, significant declines in many areas of the state have been documented (see Williams and others, 2008, for a review). With the rapid increase in urbanization and associated pollution and uncontrolled runoff from mining, farming, and silvicultural activities leading to sharply elevated sediment loads and nutrification, many areas of the state face further declines if measures to protect the habitat of mussels and their host fishes are not established and enforced.

In 1990 the U.S. Fish and Wildlife Service assessed the mussel fauna at 73 stations in the Black Warrior River system, mostly at locations where historic records were available for comparison, but also at some stations where historic data were not available (Hartfield, 1990). Collections were made by hand while digging in the substrate for live mussels, walking stream banks and gravel bars collecting material cast up by the river, and with SCUBA gear in impounded reaches. No information on effort expended in collecting was reported for comparison. Sixteen collections were made in the Locust Fork proper and 14 other collections among Blackburn Fork, Little Warrior River, and Gurley, Mud, Turkey and Valley Creeks. The best habitat in the main channel was between Alabama Highway 77 near Warrior downstream to the mouth of Shoal Creek. No mussels were found upstream of the confluence of Blackburn Fork. Only 2 species were collected live from the main channel Locust Fork, 3 fresh dead, and 12 weathered dead or relict material only. The Locust Fork was silted in the upper reaches, likely due to truck farms, irrigation, chicken farming, and coal strip mining. Among the tributary collections only a single weathered dead shell was found in Gurley Creek and a relict shell in Blackburn Fork.

The purpose of the present study is twofold: first, to document current populations of federally listed and State of Alabama Conservation Priority mussel species (Mirarchi, 2004) at selected stations in the Black Warrior River system; and second, to document other species in the system. It is hoped that information gathered during this study will serve as a guide for regulatory agencies to target streams in need of protection and provide stakeholders with

information to lead them in appropriate activities to protect the existing fauna and foster recovery of the fauna as a whole.

ACKNOWLEDGMENTS

The authors gratefully acknowledge Rong Li of the University of Alabama Department of Information Systems, Statistics, and Management Science, Prarthana Ghosh of the University of Alabama Department of Biological Sciences, Cal Johnson, Pat O'Neil, Brett Smith, and Tom Shepard of the Geological Survey of Alabama, and Patti Pennington and Matthew Nance of Cahaba/Black Warrior/Coosa Resource Conservation & Development Council for assistance with field sampling. Funding for this project was provided by the Alabama Department of Conservation and Natural Resources through Section 6 of the Endangered Species Act.

STUDY AREA

The Mobile River Basin is the largest Gulf of Mexico river basin east of the Mississippi River, draining about 43,683 square miles (mi²) in Alabama, Mississippi, Georgia, and Tennessee, including 32,207 mi², or 62 percent, of the land area of Alabama (Mettee and others, 1996). The Tombigbee River system is the westernmost tributary of the Mobile Basin and drains an area of 19,984 mi² in Alabama and Mississippi, and the Black Warrior River drains 6,228 mi² in north-central Alabama (fig. 1). Sampling in the main channel Black Warrior River extended from the tailwater of Oliver Lock and Dam at Black Warrior River mile (BWM) 113.6 upstream in the Oliver and Holt Pools to the Bankhead Lock and Dam tailwater at BWM 146.6. Tributary stations were all located upstream of the Fall Line and were selected based on accessibility and presence of suitable habitat for target species. Sampling locations were determined with the aid of Black Warrior River charts, DeLorme's Atlas and Gazetteer, and hand-held Global Positioning System (GPS) units.

METHODS

Mussel sampling was performed at stations within the Black Warrior River system where mussels were known historically and in streams with no historic data available. Stations were accessed at bridge crossings, foot trails, or by boat. Mussel collections were made by hand often with the aid of mask and snorkel in tributary stations and with a surface air source at main channel stations. Sampling in tributaries was restricted to larger tributaries of the Black Warrior River upstream of the Fall Line, an area where numerous species, including current federally listed and State Conservation Priority species, were historically reported. Main channel stations




Figure 1. Map of the study area showing mussel sampling locations.

were randomly selected on presumed presence of suitable habitat for mussels from the tailwater of Oliver Lock and Dam near Tuscaloosa upstream to the tailwater of Bankhead Lock and Dam in the Holt Pool.

Due to the nature of the project and limited sampling time and resources, a generally qualitative sampling protocol (timed search) was employed, with emphasis on sampling habitats favored by target species (Strayer and Smith, 2003). Sampling time was dictated by the habitat or fauna encountered at each station. If a meager fauna or poor quality habitat was encountered, we terminated our efforts at that station and moved to another sampling station. Stations where the potential for finding target species was highest (generally stable substrate with mixed sand, gravel, cobble, or boulders) received more attention; therefore, species that prefer other habitats may be underrepresented. Some collections were made incidental to sampling for unrelated projects and did not involve concerted effort to find live animals.

Live animals were identified and returned to the stream where they were found. A few problematic specimens were retained for verification and possible genetic work and were deposited in the University of Alabama Malacological Collection. Representative specimens of shell material collected were retained and will be deposited in the North Carolina State Museum of Natural Sciences or the University of Alabama Malacological Collection. Nomenclature follows Williams and others (2008) and notes on species distributions were gleaned from Williams and others (2008) and Mirarchi (2004).

RESULTS AND DISCUSSION

During this project 16 stations were sampled in the main channel Black Warrior River and 43 in tributaries (table 1, fig. 1). Approximately 8 hours of bottom time were spent sampling in the main channel (average 0.5 hour per station) and 58 hours in tributaries (average 0.7 hour per station). An aggregate total of 26 species were collected, with 16 found in the main channel and 17 in tributaries, with 16 species common to both habitats (tables 2, 3).

A cumulative total of 675 mussels either live or fresh dead were collected in main channel stations, for a catch per unit effort (CPUE) of 84.4 mussels/hour. The most numerically dominant and frequently encountered species in the main channel were the Bankclimber, *Plectomerus dombeyanus* (9 stations, CPUE 28.4), the Southern Mapleleaf, *Quadrula apiculata* (11 stations, CPUE 23.8), and the Threehorn Wartyback, *Obliquaria reflexa* (11 stations, CPUE 22.0). Only four individuals of the federally threatened Inflated Heelsplitter, *Potamilus inflatus*,

Table 1-Summary information for stations sampled in the Black Warrior River system, Alabama, 2009-10.

Locality ¹	County	Map coordinates
Main channel stations¹		
BWM 113.6 opposite Snow's Drift in Clement Bend, Oliver tailwater	Tuscaloosa	N 33.1954° W 87.6794°
BWM 116.5, Oliver tailwater near mouth of Big Creek	Tuscaloosa	N 33.2028° W 87.6605°
BWM 118.5 downstream of Black Warrior Parkway, Oliver tailwater	Tuscaloosa	N 33.1972° W 87.6286°
BWM 121.3 near old Oliver Lock and Dam, Oliver Pool	Tuscaloosa	N 33.2113° W 87.5821°
BWM 122.0 at Bama Belle mooring facility, Oliver Pool	Tuscaloosa	N 33.2141° W 87.5723°
BWM 123.4 downstream of old lock wall near Tuscaloosa, Oliver Pool	Tuscaloosa	N 33.2304° W 87.5500°
BWM 123.6 upstream of old lock wall near Tuscaloosa, Oliver Pool	Tuscaloosa	N 33.2221° W 87.5459°
BWM 124.8, Oliver Pool upstream of U.S. Highway 82 bridge	Tuscaloosa	N 33.2241° W 87.5246°
BWM 126.3 downstream of new bridge, Oliver Pool	Tuscaloosa	N 33.2395° W 87.5083°
BWM 128.0, Oliver Pool near Waterfalls Branch	Tuscaloosa	N 33.2524° W 87.4808°
BWM 129.1 upstream of Hurricane Creek mouth, Oliver Pool	Tuscaloosa	N 33.2529° W 87.4648°
BWM 133.3 upstream of Rocky Branch boat ramp, Holt Pool	Tuscaloosa	N 33.2811° W 87.4214°
BWM 137.5 upstream of Bluff Creek at Laurel Branch, Holt Pool	Tuscaloosa	N 33.3184° W 87.4140°
BWM 141.4 downstream of Harold's Lake, Holt Pool	Tuscaloosa	N 33.3715° W 87.4095°
BWM 144.4 opposite mouth of Davis Creek, Holt Pool	Tuscaloosa	N 33.4090° W 87.3945°
BWM 146.6 at mouth of Blue Creek, Holt Pool/Bankhead tailwater	Tuscaloosa	N 33.4365° W 87.3796°
Black Warrior River tributary stations		
Davis Creek at County Road 59 near Kellerman	Tuscaloosa	N 33.3882° W 87.2969°
Davis Creek at Hannah Road	Tuscaloosa	N 33.3319° W 87.2375°
Davis Creek downstream of County Road 99 near Patterson town	Tuscaloosa	N 33.3104° W 87.2221°
Blue Creek at Alabama Highway 69	Tuscaloosa	N 33.5218° W 87.4849°
Big Yellow Creek at Alabama Highway 69	Tuscaloosa	N 33.5684° W 87.4080°
Wolf Creek at Alabama Highway 18 W of Oakman	Walker	N 33.7088° W 87.4777°
Wolf Creek alongside County Road 173	Walker	N 33.7300° W 87.4724°
Wolf Creek at Wolf Creek Road upstream of Alabama Highway 102	Fayette	N 33.7994° W 87.5334°
Lost Creek at County Road 20 W of Parrish and E of Oakman	Walker	N 33.7250° W 87.3111°
Lost Creek at Pleasant Grove Road SW of New Jagger	Walker	N 33.8025° W 87.3679°
Lost Creek at Alabama Highway 118 near Carbon Hill	Walker	N 33.8816° W 87.5098°
Mill Creek at Radiant City Road near Spring Hill, 2 mi SW of Nauvoo	Walker	N 33.9705° W 87.5378°
Lost Creek at Haley Bottoms Road near Kansas community	Walker	N 33.9019° W 87.5710°
Locust Fork and tributary stations		
Valley Creek near Oak Grove at Lock 17 Road (Co. Road 54)	Jefferson	N 33.4469° W 87.1225°
Village Creek at Elbow Porter Road near West Jefferson Power Plant	Jefferson	N 33.6274° W 87.0531°
Village Creek at Mulga Mine near Maytown	Jefferson	N 33.5666° W 87.0025°
Locust Fork at shoal 1 mile downstream of railroad bridge	Jefferson	N 33.6792° W 87.0026°
Five Mile Creek at U.S. Highway 78 near Graysville	Jefferson	N 33.6636° W 86.9709°

Table 1-Summary information for stations sampled in the
Black Warrior River system, Alabama, 2009-10—continued.

Locality ¹	County	Map coordinates
Five Mile Creek at Brookside Road near Brookside	Jefferson	N 33.6398° W 86.9161°
Five Mile Creek at Upper Coalburg on Coalburg Road	Jefferson	N 33.6060° W 86.8541°
Ward Creek at County Road 140 (Warrior-Jasper Road)	Jefferson	N 33.7641° W 86.9249°
Turkey Creek at Morris-Majestic Road 0.5 mile W of Crosston	Jefferson	N 33.7292° W 86.7391°
Turkey Creek at Turkey Creek Road in Turkey Creek Nature Preserve	Jefferson	N 33.7046° W 86.6948°
Gurley Creek at County Road 133 near Trafford	Jefferson	N 33.8023° W 86.7532°
Longs Branch at County Road 22 near County Line community	Blount	N 33.8388° W 86.7272°
Locust Fork at Dean's Ferry Bridge on County Road 22	Blount	N 33.8463° W 86.7274°
Locust Fork at County Road 43, Vaughn's Bridge, on Wallston Road	Blount	N 33.8893° W 86.6955°
Locust Fork at County Road 13	Blount	N 33.9459° W 86.6691°
Locust Fork at shoal 0.5 mi downstream of Blackburn Fork confluence	Blount	N 33.9354° W 86.6442°
Blackburn Fork at low water bridge on Jerry Marsh Road	Blount	N 33.9351° W 86.6159°
Calvert Prong at Moss Bridge on Deavers Town Road	Blount	N 33.3508° W 86.5825°
Calvert Prong at County Road 33	Blount	N 33.9773° W 86.5272°
Blackburn Fork at Hendrick Mill on House Road off County Road 15	Blount	N 33.8798° W 86.5805°
Hendrick Mill Branch at County Road 15 near Limestone Springs	Blount	N 33.8771° W 86.5677°
Blackburn Fork at Alabama Highway 75 near Remlap	Blount	N 33.8566° W 86.5632°
Blackburn Fork at Blount County Road 27	Blount	N 33.8654° W 86.4435°
Blackburn Fork at Blount County Road 20	Blount	N 33.9059° W 86.3934°
Locust Fork at Alabama Highway 160 near Nectar	Blount	N 33.9803° W 86.6159°
Slab Creek at County Road 39	Marshall	N 34.1948° W 86.3619°
Big Mud Creek at County Road 21	Blount	N 34.1354° W 86.3720°
Locust Fork at unnumbered county road 1 mile NE of Walnut Grove	Etowah	N 34.0844° W 86.2892°
Bristow Creek at Bud Umphrey Road near Pine Grove	Etowah	N 34.0881° W 86.2573°
Locust Fork at Dee Nix Road near Altoona	Etowah	N 33.9956° W 86.3103°

¹ BWM = Black Warrior River mile

Table 2.—Overview of freshwater mussels collected in the Black Warrior River, Alabama, 2009.

Species	Status ¹	Results of sampling, summer 2009
<i>Anodonta suborbiculata</i> , Flat Floater	P4	Two live individuals were found in the upper reach of Holt Pool; these are the first main channel Black Warrior records of this adventitious species
<i>Arcidens confragosus</i> , Rock Pocketbook	P3	Two live individuals were found in the Oliver Pool near Tuscaloosa
<i>Lampsilis teres</i> , Yellow Sandshell	P5	Live individuals were frequently encountered throughout the study area
<i>Lasmigona alabamensis</i> , Alabama Heelsplitter	P3	Live individuals were frequently encountered throughout the study area
<i>Leptodea fragilis</i> , Fragile Papershell	P5	Live individuals were frequently encountered throughout the study area
<i>Megalonaias nervosa</i> , Washboard	P5	A few live individuals were found at two stations in the upper reach of Holt Pool
<i>Obliquaria reflexa</i> , Threehorn Wartyback	P5	The third most frequently encountered and numerically abundant species encountered, it was common throughout the study area
<i>Plectomerus dombeyanus</i> , Bankclimber	P5	The most frequently encountered and numerically abundant species encountered, it was common throughout the study area
<i>Potamilus inflatus</i> , Inflated Heelsplitter	T, P2	Four live individuals were encountered at two stations, one downstream of Oliver Dam and one in the Oliver Pool near Tuscaloosa
<i>Potamilus purpuratus</i> , Bleufer	P5	Live individuals were frequently encountered throughout the study area
<i>Pyganodon grandis</i> , Giant Floater	P5	A few live individuals were found at several scattered stations in the study area
<i>Quadrula apiculata</i> , Southern Mapleleaf	P5	The second most frequently encountered and numerically abundant species encountered, it was common throughout the study area
<i>Quadrula asperata</i> , Alabama Orb	P5	Only one live individual of this otherwise common and widespread Mobile Basin endemic was encountered in the Oliver Pool near Tuscaloosa
<i>Quadrula rumphiana</i> , Ridged Mapleleaf	P4	A fairly commonly encountered species, especially in the Holt Pool; it can be difficult to distinguish from <i>Quadrula apiculata</i> , with some evidence of intergradation
<i>Toxolasma parvum</i> , Lilliput	P3	A single live individual of this diminutive species was found in the Holt Pool
<i>Utterbackia imbecillis</i> , Paper Pondshell	P5	A few live individuals and fresh dead shells of this common and widespread species were found

¹ T=federally listed threatened; Alabama priority conservation ranks follow Mirarchi (2004): P2=High Conservation Concern, P3=Moderate Conservation Concern, P4=Low Conservation Concern, P5=Lowest Conservation Concern.

Table 3. — Overview of freshwater mussels collected in tributaries of the Black Warrior River, Alabama, 2009-10.

Species	Status ¹	Results of sampling, summer 2009 and 10
<i>Amblema plicata</i> , Threeridge	P5	Weathered dead valves were found in Locust Fork at Dean's Ferry Bridge and at Blount County Road 13
<i>Elliptio arctata</i> , Delicate Spike	P1	Two weathered dead valves were found in Locust Fork at Vaughn's Bridge
<i>Lampsilis ornata</i> , Southern Pocketbook	P4	Two live animals (including one female displaying) were found at the lowermost station in Blackburn Fork; numerous live animals were found in Locust Fork at Dean's Ferry Bridge and Vaughn's Bridge while dead shells were found at several nearby stations
<i>Lampsilis straminea</i> , Southern Fatmucket	P4	One valve of a relic shell was found in Turkey Creek
<i>Lampsilis teres</i> , Yellow Sandshell	P5	Live animals were found in Locust Fork at Dean's Ferry Bridge and Vaughn's Bridge; a weathered dead shell was found in Village Creek at the lowermost station
<i>Lasmigona alabamensis</i> , Alabama Heelsplitter	P3	Single fresh dead shells were found in Locust Fork at Vaughn's Bridge and at Blount County Road 13, and in Village Creek at the lowermost station
<i>Leptodea fragilis</i> , Fragile Papershell	P5	Several weathered dead valves were found in Village Creek at the two lowermost stations and in Locust Fork at Vaughn's Bridge and Blount County Road 13
<i>Obliquaria reflexa</i> , Threehorn Wartyback	P5	A few live animals were found in Locust Fork at Dean's Ferry Bridge and Vaughn's Bridge and weathered dead at Blount County Road 13
<i>Pleurobema rubellum</i> , Warrior Pigtoe	X	One valve of a relic shell tentatively identified as this species was collected from Davis Creek at Hannah Road
<i>Potamilus purpuratus</i> , Bleufer	P5	A live animal was found in Locust Fork at County Road 13 and some fresh and weathered dead shells were found at several nearby stations; 25 live animals were found at the lowermost station in Blackburn Fork; fresh dead and weathered dead shells were found at the two lowermost stations in Village Creek
<i>Pyganodon grandis</i> , Giant Floater	P5	Three weathered dead shells were found in Village Creek at the lowermost station

Table 3. — Overview of freshwater mussels collected in tributaries of the Black Warrior River, Alabama, 2009-10—continued.

Species	Status ¹	Results of sampling, summer 2009 and 10
<i>Quadrula asperata</i> , Alabama Orb	P5	Single weathered dead valves were found at each of the lowermost Davis Creek stations; live animals were found in Locust Fork at Dean's Ferry Bridge and Vaughn's Bridge
<i>Quadrula verrucosa</i> , Pistolgrip	P4	Three live animals were found at the lowermost station in Blackburn Fork; in Locust Fork a few live animals were found downstream of the mouth of Blackburn Fork and at Blount County Road 13 and fresh dead shells were found at Vaughn's Bridge
<i>Strophitus subvexus</i> , Southern Creekmussel	P3	One partial valve of a relic shell was found at the Blue Creek station
Unionidae	X	One badly weathered valve of a Unionid, probably <i>Lampsilis</i> sp., was found in Five Mile Creek at Brookside
<i>Villosa lienosa</i> , Little Spectaclecase	P5	One fresh dead and two weathered dead shells were found at the Blue Creek station
<i>Villosa nebulosa</i> , Alabama Rainbow	P3	Two fresh dead shells were found at the Blue Creek station and two weathered dead shells in Blackburn Fork
<i>Villosa vibex</i> , Southern Rainbow	P5	Two weathered dead shells were found in Blackburn Fork at County Road 27

¹ Alabama priority conservation ranks follow Mirarchi (2004): P1=Highest Conservation Concern, P3=Moderate Conservation Concern, P4=Low Conservation Concern, P5=Lowest Conservation Concern, X=not assigned.

were collected, with three live individuals collected in the Oliver Lock and Dam tailwater at BWM 118.5 near the Black Warrior Parkway Bridge, and one live individual near the city of Tuscaloosa at BWM 122.0. A breakdown of species encountered is presented in table 2.

Mussels in the main channel Black Warrior were usually found in areas of stable gravel and sand sometimes mixed with cobble and boulders and generally having varying levels of silt deposition. The two most downstream main channel collections (BWM 113.6 and 116.5, table 1, fig. 1), located in the Oliver Lock and Dam tailwater just downstream of the Fall Line on the Coastal Plain, had relatively poor habitat and little effort was spent sampling there. Habitat generally improved with upstream progression in the main channel. Habitat in the Oliver and Holt pools was comprised primarily of fairly stable gravel and sand substrates with occasional cobble, boulders, woody debris, and bedrock with some areas of mud, often with a layer of fine silt in eddies and areas protected from the current. Visibility was often 3 to 4 feet with a light source.

Over the two summers of this study 30 stations were sampled in the Locust Fork system, with 8 in the main channel and 22 in selected tributaries (fig. 1, table 1). Hartfield (1990) reported on species collected from 16 main channel Locust Fork stations and 14 additional stations among Gurley, Mud, Turkey, and Valley Creeks, Blackburn Fork, and Little Warrior River. During the present study the main channel Locust Fork and the lower reach of its major tributary, Blackburn Fork, yielded relatively diverse populations of fairly common mussels (seven species) as well as a few old shells of more sensitive species, providing hope that with further exploration extant populations of rare species may be found (table 4). Most tributaries yielded few or no mussels. Hartfield (1990) reported two live, three fresh dead, and 12 weathered dead or relict shells from main channel Locust Fork stations, a single weathered dead shell of *Quadrula verrucosa* in Gurley Creek, and a relict shell of *Lampsilis teres* in Blackburn Fork. No mussels were found in the Locust Fork main channel upstream of the confluence of Blackburn Fork.

A cumulative total of 13 species was found in Locust Fork and tributaries during the present study, compared to 17 reported by Hartfield (1990) (total 21 between the two studies). A comparison of the results of the respective studies is summarized in table 4. Nine species were common to both studies. Species reported by Hartfield (1990) not found during the current study include *Elliptio arca*, *Elliptio crassidens*, *Fusconaia cerina*, *Ligumia recta*, *Megaloniaias*

Table 4. — Comparison of freshwater mussel species in Locust Fork and tributaries reported by Hartfield (1990) and the current study.

Species	Status ¹	Conditions of species reported ²	
		Hartfield (1990)	Current study
<i>Amblema plicata</i> , Threeridge	P5	--	W
<i>Ellipsaria lineolata</i> , Butterfly	P4	W	F
<i>Elliptio arca</i> , Alabama Spike	P5	W	--
<i>Elliptio arctata</i> , Delicate Spike	P1	W	W
<i>Elliptio crassidens</i> , Elephantear	P1	W	--
<i>Fusconaia cerina</i> , Gulf Pigtoe	P5	W	--
<i>Lampsilis ornata</i> , Southern Pocketbook	P4	L	L
<i>Lampsilis teres</i> , Yellow Sandshell	P5	W	L
<i>Lasmigona alabamensis</i> , Alabama Heelsplitter	P3	W	F
<i>Leptodea fragilis</i> , Fragile Papershell	P5	L	F
<i>Ligumia recta</i> , Black Sandshell	P4	W	--
<i>Megaloniaias nervosa</i> , Washboard	P5	W	--
<i>Obliquaria reflexa</i> , Threehorn Wartyback	P5	--	L
<i>Obovaria</i> sp.	--	W	--
<i>Pleurobema furvum</i> , Dark Pigtoe	P5	W	--
<i>Potamilus purpuratus</i> , Bleufer	P5	F	L
<i>Quadrula asperata</i> , Alabama Orb	P5	W	L
<i>Quadrula rumphiana</i> , Ridged Mapleleaf	P4	F	--
<i>Quadrula verrucosa</i> , Pistolgrip	P4	F	L
<i>Villosa nebulosa</i> , Alabama Rainbow	P3	--	W
<i>Villosa vibex</i> , Southern Rainbow	P5	--	W

¹ Alabama priority conservation ranks follow Mirarchi (2004):P1=Highest Conservation Concern, P3=Moderate Conservation Concern, P4=Low Conservation Concern, P5=Lowest Conservation Concern.

²F=fresh dead shells, L=live animals, W=weathered dead or relict shells, --=not collected.

nervosa, *Pleurobema furvum*, *Quadrula rumphiana*, and *Obovaria* sp. Further sampling at additional stations would likely add some of these species, which are generally found in larger streams than those sampled during this study. Additionally, a single fresh dead specimen of *Ptychobranthus greenii*, the Triangular Kidneyshell, a federally listed endangered species and Conservation Priority 1 species in Alabama, was collected in the main channel Locust Fork by Geological Survey of Alabama personnel in the mid-90s during an unrelated project.

SPECIES ACCOUNTS

Amblema plicata, Threeridge, P4, is a widespread and common species in the Tennessee River system, the western and lower Mobile River Basin, and some coastal river systems, and can be found in riverine or impounded areas. It is declining in the Mobile River Basin. It was found as weathered dead material at two main channel Locust Fork stations.

Anodonta suborbiculata, Flat Floater, P4, is widespread in the Tennessee River system with localized populations in Mobile River Basin impoundments and is usually found in soft sediments in sluggish water. It may be a relatively recent invader of the Mobile River Basin. Two live animals were found in the upper reach of Holt Reservoir, representing the first main channel Black Warrior River record for this species.

Arcidens confragosus, Rock Pocketbook, P3, is fairly common in some areas of the Tennessee River system but is declining in the Mobile River Basin and can be found in either riverine or impounded areas. Two live individuals were found in the Oliver Pool.

Elliptio arctata, Delicate Spike, P1, is widespread but never common in the Mobile River Basin and is declining and is almost always found in riverine habitats, often under large rocks. Two weathered dead valves were found in the Locust Fork main channel. Hartfield (1990) reported weathered dead or relict shells from Locust Fork.

Lampsilis ornata, Southern Pocketbook, P4, is widespread and common in the Mobile River Basin and sporadically in the Conecuh River drainage and can occupy a variety of habitats. Live animals (including one female displaying) were found in Locust Fork and Blackburn Fork. Hartfield (1990) reported four live individuals from Locust Fork.

Lampsilis straminea, Southern Fatmucket, P4, is fairly common and widespread in Alabama south of the Tennessee River system and is often found in slow to moderate current but generally not in impoundments. A relict shell was found in Turkey Creek.

Lampsilis teres, Yellow Sandshell, P5, is common throughout Alabama and may be found in gravel, sand, or mud substrates in riverine or impounded habitats. It was frequently encountered in the Black Warrior main channel and Locust Fork and a weathered dead shell was found in Village Creek. Hartfield (1990) reported weathered dead or relict shells from Locust Fork.

Lasmigona alabamensis, Alabama Heelsplitter, P3, is restricted to the Mobile River Basin and is uncommon and may be found in riverine or pooled habitats. It was found live at numerous stations in the main channel Black Warrior River and fresh dead at several stations in Locust Fork and one station in Village Creek. Hartfield (1990) reported weathered dead or relict shells from Locust Fork.

Leptodea fragilis, Fragile Papershell, P5, is a common and widespread species in the Tennessee River system and Mobile River Basin and can be found in riverine habitats and impoundments. It was frequently encountered live in the main channel Black Warrior and as weathered dead valves in Locust Fork and Village Creek. Hartfield (1990) reported a live individual from Locust Fork.

Megaloniais nervosa, Washboard, P5, is common and widespread throughout Alabama except the Choctawhatchee and Yellow River systems, and can be found in riverine habitats and impoundments. It was found live at a few stations in the upper reach of Holt Pool. Hartfield (1990) reported weathered dead or relict shells from Locust Fork.

Obliquaria reflexa, Threehorn Wartyback, P5, is common throughout the Tennessee River system and Mobile River Basin, and can be found in riverine habitats and impoundments. It was frequently encountered at main channel Black Warrior River stations and was the third most frequently collected species there. A few live animals were found in the Locust Fork.

Plectomerus dombeyanus, Bankclimber, P5, is common in the Alabama and lower Tombigbee River drainages and lower reaches of Coosa and Cahaba Rivers, and occurs in both sluggish and flowing water, often on channel slopes. It was the most commonly encountered and numerically abundant species found in the main channel Black Warrior River.

Pleurobema rubellum, Warrior Pigtoe, considered extinct, was endemic to the Black Warrior and Cahaba River systems, probably in stable gravel substrates like its congeners, but has not been collected in nearly 100 years. One very old shell resembling this species was found in Davis Creek.

Potamilus inflatus, Inflated Heelsplitter, P2, a federally listed threatened species, is uncommon, restricted to the Mobile River Basin, and is usually found in soft substrates in slow to moderate current. Four live individuals were collected in the Black Warrior River: three in the tailwater of Oliver Dam and one in the Oliver Pool near Tuscaloosa.

Potamilus purpuratus, Bleufer, P5, is widespread and common in the Mobile River Basin and is found in both pools and shoals, often under large rocks in shoals. It was found frequently among the main channel Black Warrior River stations. One live individual and several fresh dead shells were found among several Locust Fork stations, and 25 live animals were found at the most downstream station in Blackburn Fork. Hartfield (1990) reported fresh dead shells from Locust Fork.

Pyganodon grandis, Giant Floater, P5, is common throughout Alabama and occurs in practically any habitat including farm ponds. Live animals were found at several stations in the main channel Black Warrior River and weathered dead shells were found in Village Creek.

Quadrula apiculata, Southern Mapleleaf, P5, is common and was formerly restricted to the Mobile River Basin but has been introduced into the Tennessee River, likely by mussel divers, and can be found in riverine and impounded reaches. It was the second most commonly encountered and numerically abundant species found among Black Warrior River stations.

Quadrula asperata, Alabama Orb, P5, is common and endemic to the Mobile River Basin, usually in habitats with at least some current. Oddly, only one live individual of this common and widespread Mobile River Basin endemic was found in the Oliver Pool of Black Warrior River. Live animals were found at two stations in Locust Fork and a few weathered dead valves were found in Davis Creek. Hartfield (1990) reported weathered dead or relict shells from Locust Fork.

Quadrula rumphiana, Ridged Mapleleaf, P4, is endemic to the Mobile River Basin and fairly common, usually in habitats with at least some current. It was commonly encountered in main channel Black Warrior River stations, especially in the Holt Pool, but not in tributaries. Hartfield (1990) reported fresh dead shells from Locust Fork.

Quadrula verrucosa, Pistolgrip, P4, is fairly common and widespread in the Tennessee River system and Mobile River Basin, usually in streams with at least some current but occasionally in impoundments. Live animals were found in lower Blackburn Fork and at several stations in Locust Fork. Hartfield (1990) reported fresh dead shells from Locust Fork.

Strophitus subvexus, Southern Creekmussel, P3, is found throughout Alabama south of the Tennessee River system but is uncommon and is often found in sluggish streams. One relict valve was found in Blue Creek.

Toxolasma parvum, Lilliput, P3, is known from the Tennessee River system, Mobile River Basin, and Gulf Coast drainages, and often occupies soft sediments in sluggish water. It is poorly known and further taxonomic work may lead to generic revision and further restrict its distribution. A single live individual was found in the Holt Pool of the Black Warrior River.

Utterbackia imbecillis, Paper Pondshell, P5, is common throughout Alabama and occurs in practically any habitat including farm ponds. A few live and fresh dead individuals were found in the main channel Black Warrior River.

Villosa lienosa, Little Spectaclecase, P5, is common throughout Alabama south of the Tennessee River system and can occur in a variety of habitats. One fresh dead and a few weathered dead shells were found in Blue Creek.

Villosa nebulosa, Alabama Rainbow, P3, is occasionally found in the Mobile River Basin upstream of the Fall Line, usually in small stream environments. Two fresh dead shells were found in Blue Creek and two weathered dead shells in Blackburn Fork.

Villosa vibex, Southern Rainbow, P5, is common throughout Alabama south of the Tennessee River system in a variety of habitats. Two weathered dead shells were found in Blackburn Fork.

RECOMMENDATIONS

Based on the results of this study we make the following recommendations:

- Further sampling of the mussel fauna should be executed to further refine the current distribution and status of mussels in the system.
- Factors that influence mussel distribution and abundance should be evaluated by such means as land cover/land use determinations, water and sediment quality measurements, and evaluation of rates of sediment loading.
- Upon determination of the factors limiting the fauna, steps should be taken to ameliorate those factors.
- Periodic monitoring of selected stations should be executed to document trends.

REFERENCES CITED

- Hartfield, Paul, 1990, Status survey for mussels in the tributaries of the Black Warrior River, Alabama, 1990: Unpublished report from U.S. Fish and Wildlife Service, Jackson, Mississippi, 8 p.
- Hartfield, Paul, 1994, Status review of select mussel species in the Mobile River basin: U.S. Fish and Wildlife Service, Jackson, Mississippi, 3 p.
- Hinkley, A. A., 1906, Some shells from Mississippi and Alabama: *The Nautilus*, v. 20, nos. 3, 4, and 5, p. 34-36, 40-44, and 52-55.
- Mettee, M. F., O'Neil, P. E., and Pierson, J. M., 1996, Fishes of Alabama and the Mobile basin including Mobile basin tributaries in adjacent states: Oxmoor House, Birmingham, Alabama, 820 p.
- Mirarchi, R. E., 2004, Alabama wildlife, Volume 1, A checklist of vertebrates and selected invertebrates: Aquatic mollusks, fishes, amphibians, reptiles, birds, and mammals: Tuscaloosa, The University of Alabama Press, 209 p.
- Mott, Seth, and Hartfield, Paul, 1994, Status review summary of the Alabama pearlshell, *Margaritifera marrianae*: U.S. Fish and Wildlife Service, Jackson, Mississippi, 6 p.
- Strayer, D. L., and Smith, D. R., 2003, A guide to sampling freshwater mussel populations: Bethesda, Maryland, American Fisheries Society, Monograph 8, 103 p.
- Williams, J. D., 1982, Distribution and habitat observations of selected Mobile basin unionid mollusks, in Miller, A. C., ed., Report of freshwater mollusks workshop: Vicksburg, Mississippi, U.S. Army Corps of Engineers, Waterways Experiment Station, p. 61-85.
- Williams, J.D., Bogan, A.E., and Garner, J.T., 2008, Freshwater mussels of Alabama and the Mobile basin in Georgia, Mississippi, and Tennessee: The University of Alabama Press, Tuscaloosa, 908 p.
- Williams, J. D., Fuller, S. L., and Grace, Randall, 1992, Effects of impoundments on freshwater mussels (Mollusca: Bivalvia: Unionidae) in the main channel of the Black Warrior and Tombigbee Rivers in western Alabama: *Bulletin of the Alabama Museum of Natural History*, v. 13, p. 1-10.

GEOLOGICAL SURVEY OF ALABAMA

P.O. Box 869999
420 Hackberry Lane
Tuscaloosa, Alabama 35486-6999
205/349-2852

Berry H. (Nick) Tew, Jr., State Geologist

A list of the printed publications by the Geological Survey of Alabama can be obtained from the Publications Sales Office (205/247-3636) or through our web site at <http://www.gsa.state.al.us/>.

E-mail: pubsales@gsa.state.al.us

The Geological Survey of Alabama (GSA) makes every effort to collect, provide, and maintain accurate and complete information. However, data acquisition and research are ongoing activities of GSA, and interpretations may be revised as new data are acquired. Therefore, all information made available to the public by GSA should be viewed in that context. Neither the GSA nor any employee thereof makes any warranty, expressed or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed in this report. Conclusions drawn or actions taken on the basis of these data and information are the sole responsibility of the user.

As a recipient of Federal financial assistance from the U.S. Department of the Interior, the GSA prohibits discrimination on the basis of race, color, national origin, age, or disability in its programs or activities. Discrimination on the basis of sex is prohibited in federally assisted GSA education programs. If anyone believes that he or she has been discriminated against in any of the GSA's programs or activities, including its employment practices, the individual may contact the U.S. Geological Survey, U.S. Department of the Interior, Washington, D.C. 20240.

AN EQUAL OPPORTUNITY EMPLOYER

Serving Alabama since 1848